

## New England Biolabs Certificate of Analysis

**Product Name:** Pyrophosphatase, Inorganic (*E. coli*)  
**Catalog Number:** M0361L  
**Concentration:** 100 U/ml  
**Unit Definition:** One unit is the amount of enzyme that will generate 1  $\mu$ mol of phosphate per minute from inorganic pyrophosphate under standard reaction conditions.  
**Lot Number:** 10043148  
**Expiration Date:** 04/2021  
**Storage Temperature:** -20°C  
**Storage Conditions:** 100 mM NaCl, 20 mM Tris-HCl (pH 8.0), 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol  
**Specification Version:** PS-M0361S/L v2.0


### Pyrophosphatase, Inorganic (*E. coli*) Component List

NEB Part Number	Component Description	Lot Number	Individual QC Result
M0361LVIAL	Pyrophosphatase, Inorganic ( <i>E. coli</i> )	10043149	Pass


Assay Name/Specification	Lot # 10043148
<b>Endonuclease Activity (Nicking)</b> A 50 $\mu$ l reaction in NEBuffer 4 containing 1 $\mu$ g of supercoiled PhiX174 DNA and a minimum of 0.5 units of Pyrophosphatase, Inorganic ( <i>E. coli</i> ) incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 $\mu$ l reaction in NEBuffer 4 containing 1 $\mu$ g of a mixture of single and double-stranded [ <sup>3</sup> H] <i>E. coli</i> DNA and a minimum of 0.5 units of Pyrophosphatase, Inorganic ( <i>E. coli</i> ) incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Phosphatase Activity (pNPP)</b> A 100 $\mu$ l reaction in NEBuffer 4 containing 10 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 1 unit Pyrophosphatase, Inorganic ( <i>E. coli</i> ) incubated for 1 hour at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass
<b>Protein Purity Assay (SDS-PAGE)</b>	Pass

Assay Name/Specification	Lot # 10043148
<p>Pyrophosphatase, Inorganic (E. coli) is <math>\geq 95\%</math> pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p> <p><b>RNase Activity (Extended Digestion)</b> A 10 <math>\mu</math>l reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 0.1 unit of Pyrophosphatase, Inorganic (E. coli) is incubated at 37°C. After incubation for 16 hours, &gt;90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	<p><b>Pass</b></p>

This product has been tested and shown to be in compliance with all specifications.



Dongxian Yue  
Production Scientist  
17 May 2019



Michael Tonello  
Packaging Quality Control Inspector  
22 May 2019