

## New England Biolabs Certificate of Analysis

**Product Name:** OneTaq® Hot Start 2X Master Mix with Standard Buffer  
**Catalog Number:** M0484L  
**Concentration:** 2 X Concentrate  
**Lot Number:** 10044962  
**Expiration Date:** 01/2021  
**Storage Temperature:** -20°C  
**Specification Version:** PS-M0484S/L v1.0  
**Composition (1X):** 20 mM Tris-HCl (pH 8.9 @ 25°C), 22 mM NH<sub>4</sub>Cl, 22 mM KCl, 1.8 mM MgCl<sub>2</sub>, 0.2 mM dATP, 0.2 mM dCTP, 0.2 mM dGTP, 0.2 mM dTTP, 5 % Glycerol, 0.06 % IGEPAL® CA-630, 0.05 % Tween® 20, 25 units/ml OneTaq® Hot Start DNA Polymerase

OneTaq® Hot Start 2X Master Mix with Standard Buffer Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0484SVIAL	OneTaq® Hot Start 2X Master Mix with Standard Buffer	10032983	Pass


Assay Name/Specification	Lot # 10044962
<p><b>PCR Amplification (Hot Start 2 kb Lambda DNA)</b>            A 25 µl reaction in OneTaq® Standard Reaction Buffer in the presence of 200 µM dNTPs and 0.2 µM primers containing 10 pg Lambda DNA and 50 ng Human Genomic DNA with 0.625 units of OneTaq® Hot Start DNA Polymerase for 30 cycles of PCR amplification results in an increase in yield of the 2 kb Lambda product and a decrease in non-specific genomic bands when compared to a non-hot start control reaction.</p>	Pass
<p><b>RNase Activity (Extended Digestion)</b>            A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of OneTaq® Hot Start 2X Master Mix with Standard Buffer is incubated at 37°C. After incubation for 4 hours, &gt;90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass
<p><b>Inhibition of Primer Extension (Hot Start, Radioactivity Incorporation)</b>            A 50 µl primer extension assay in ThermoPol® Reaction Buffer in the presence of 200 µM dNTPs including [ <sup>3</sup>H]-dTTP, containing 15 nM primed single-stranded M13mp18 with 2.5 units of OneTaq® Hot Start DNA Polymerase incubated for 16 hours at 25°C yields &gt;95% inhibition when compared to a non-hot start control reaction.</p>	Pass
<p><b>Non-Specific DNase Activity (16 hour, Buffer)</b></p>	Pass

Assay Name/Specification	Lot # 10044962
<p>A 50 µl reaction in 1X OneTaq<sup>®</sup> Hot Start Master Mix with Standard Buffer containing 1 µg of T3 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p> <p><b>PCR Amplification (5 kb Lambda, Master Mix)</b> A 25 µl reaction in 1X OneTaq<sup>®</sup> Hot Start Master Mix with Standard Buffer and 0.2 µM primers containing 5 ng Lambda DNA for 25 cycles of PCR amplification results in the expected 5 kb product.</p>	<p><b>Pass</b></p>

This product has been tested and shown to be in compliance with all specifications.



Christie Vazquez  
Production Scientist  
24 Jan 2019



Michael Tonello  
Packaging Quality Control Inspector  
22 May 2019