

New England Biolabs Certificate of Analysis

Product Name: Quick Ligation™ Kit
Catalog Number: M2200S
Unit Definition: N/A
Packaging Lot Number: 10236015
Expiration Date: 01/2026
Storage Temperature: -20°C
Storage Conditions: 10 mM Tris-HCl , 50 mM KCl , 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol, (pH 7.4 @ 25°C)
Specification Version: PS-M2200S/L v1.0

| Quick Ligation™ Kit Component List | | | |
|------------------------------------|---------------------------------|------------|----------------------|
| NEB Part Number | Component Description | Lot Number | Individual QC Result |
| M2200SVIAL | Quick Ligase | 10223528 | Pass |
| B2200SVIAL | Quick Ligation™ Reaction Buffer | 10220827 | Pass |

| Assay Name/Specification | Lot # 10236015 |
|--|----------------|
| DNase Activity (Labeled Oligo, 3' extension) A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a 3' extension and a minimum of 10,000 units of Quick Ligase incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis. | Pass |
| DNase Activity (Labeled Oligo, 5' extension) A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a 5' extension and a minimum of 10,000 units of Quick Ligase incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis. | Pass |
| Double Stranded DNase Activity (Labeled Oligo) A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a blunt end and a minimum of 10,000 units of Quick Ligase incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis. | Pass |
| Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 2000 units of Quick Ligase incubated for 4 hours at 37°C results in <10% | Pass |

| Assay Name/Specification | Lot # 10236015 |
|--|----------------|
| conversion to the nicked form as determined by agarose gel electrophoresis. | |
| <p>Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 1 containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 2000 units of Quick Ligase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.</p> | Pass |
| <p>Functional Testing (Ligation and Transformation) After a five-minute ligation of linearized, dephosphorylated LITMUS 28 or pUC19 (containing either blunt [EcoRV] or cohesive [HindIII] ends) and a mixture of compatible insert fragments, transformation into chemically competent E. coli DH-5 alpha cells yields a minimum of 1 x 10⁶ recombinant transformants per µg plasmid DNA.</p> | Pass |
| <p>Ligation and Recutting (Terminal Integrity, Digested DNA) A 20 µl reaction in 1X T4 DNA Ligase Reaction Buffer containing 2 µg of Lambda DNA-HindIII Digest and a minimum of 4000 units of Quick Ligase incubated for 16 hours at 37°C results in >95% ligation of the DNA fragments as determined by agarose gel electrophoresis. Of these ligated fragments, >95% can be recut with HindIII.</p> | Pass |
| <p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 1 containing 1 µg of CIP-treated Lambda-HindIII DNA and a minimum of 2000 units of Quick Ligase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p> | Pass |
| <p>Protein Purity Assay (SDS-PAGE) Quick Ligase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p> | Pass |
| <p>RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Quick Ligase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p> | Pass |
| <p>Single Stranded DNase Activity (FAM-Labeled Oligo) A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 10,000 units of Quick Ligase incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.</p> | Pass |

| Assay Name/Specification | Lot # 10236015 |
|--|--------------------|
| <p>qPCR DNA Contamination (E. coli Genomic) A minimum of 2000 units of Quick Ligase is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p> | <p>Pass</p> |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Alison Dolan
Production Scientist
07 Mar 2024



Michael Tonello
Packaging Quality Control Inspector
11 Apr 2024