

## New England Biolabs Certificate of Analysis

*Product Name:* XbaI  
*Catalog #:* R0145S/L  
*Concentration:* 20,000 units/ml  
*Unit Definition:* One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA (dam-/HindIII digest) in 1 hour at 37°C in a total reaction volume of 50 µl.  
*Lot #:* 0421504  
*Assay Date:* 04/2015  
*Expiration Date:* 04/2017  
*Storage Temp:* -20°C  
*Storage Conditions:* 50 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA  
*Specification Version:* PS-R0145S/L v1.0  
*Effective Date:* 06 Mar 2014

| Assay Name/Specification (minimum release criteria)   | Lot #0421504 |
|---|--------------|
| <b>Blue-White Screening (Terminal Integrity)</b> - A sample of pUC19 vector linearized with a 10-fold excess of XbaI, religated and transformed into an <i>E. coli</i> strain expressing the LacZ beta fragment gene results in <1% white colonies.   | <b>Pass</b>  |
| <b>Endonuclease Activity (Nicking)</b> - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 100 Units of XbaI incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.                           | <b>Pass</b>  |
| <b>Exonuclease Activity (Radioactivity Release)</b> - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] <i>E. coli</i> DNA and a minimum of 200 units of XbaI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.          | <b>Pass</b>  |
| <b>Ligation and Recutting (Terminal Integrity)</b> - After a 10-fold over-digestion of Adenovirus-2 DNA with XbaI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with XbaI.  | <b>Pass</b>  |
| <b>Non-Specific DNase Activity (16 Hour)</b> - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda HindIII dam- DNA and a minimum of 200 Units of XbaI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | <b>Pass</b>  |
| <b>Protein Purity Assay (SDS-PAGE)</b> - XbaI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.   | <b>Pass</b>  |

---


## New England Biolabs Certificate of Analysis

*\* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.*



---

Authorized by  
Derek Robinson  
06 Mar 2014



---

Inspected by  
Toby Claus  
30 Apr 2015

