

New England Biolabs Certificate of Analysis

Product Name: BclI-HF[®]
Catalog Number: R3160S
Concentration: 20,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA (dam-) in rCutSmart Buffer in 1 hour at 37°C in a total reaction volume of 50 µl.
Packaging Lot Number: 10179333
Expiration Date: 02/2025
Storage Temperature: -20°C
Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 500 µg/ml rAlbumin, (pH 7.4 @ 25°C)
Specification Version: PS-R3160S/L v2.0


| BclI-HF [®] Component List | | | |
|-------------------------------------|-------------------------------|------------|----------------------|
| NEB Part Number | Component Description | Lot Number | Individual QC Result |
| R3160SVIAL | BclI-HF [®] | 10179328 | Pass |
| B7024AVIAL | Gel Loading Dye, Purple (6X) | 10173660 | Pass |
| B6004SVIAL | rCutSmart [™] Buffer | 10175291 | Pass |

| Assay Name/Specification | Lot # 10179333 |
|--|----------------|
| Protein Purity Assay (SDS-PAGE) BclI-HF is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection. | Pass |
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in rCutSmart [™] Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 100 units of BclI-HF incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass |
| qPCR DNA Contamination (E. coli Genomic) A minimum of 20 units of BclI-HF is screened for the presence of E. coli genomic DNA using SYBR [®] Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome. | Pass |
| Functional Testing (15 minute Digest) A 50 µl reaction in rCutSmart [™] Buffer containing 1 µg of Lambda dam- DNA and 1 µl of | Pass |


| Assay Name/Specification | Lot # 10179333 |
|---|----------------|
| <p>BclI-HF incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.</p> | |
| <p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda dam- DNA and a minimum of 60 units of BclI-HF incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p> | Pass |
| <p>Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of Lambda dam- DNA with BclI-HF, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with BclI-HF.</p> | Pass |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



YunJie Sun
Production Scientist
14 Feb 2023



Michael Tonello
Packaging Quality Control Inspector
15 Feb 2023