

## New England Biolabs Certificate of Analysis

**Product Name:** *EagI-HF<sup>®</sup>*  
**Catalog Number:** *R3505S*  
**Concentration:** *20,000 U/ml*  
**Unit Definition:** *One unit is defined as the amount of enzyme required to digest 1 µg of pXba DNA in 1 hour at 37°C in a total reaction volume of 50 µl.*  
**Packaging Lot Number:** *10184482*  
**Expiration Date:** *03/2025*  
**Storage Temperature:** *-80°C*  
**Storage Conditions:** *500 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 200 µg/ml BSA, (pH 7.4 @ 25°C)*  
**Specification Version:** *PS-R3505S/L v3.0*


<b>EagI-HF<sup>®</sup> Component List</b>			
<b>NEB Part Number</b>	<b>Component Description</b>	<b>Lot Number</b>	<b>Individual QC Result</b>
R3505SVIAL	EagI-HF <sup>®</sup>	10184480	<b>Pass</b>
B7024AVIAL	Gel Loading Dye, Purple (6X)	10182168	<b>Pass</b>
B6004SVIAL	rCutSmart <sup>™</sup> Buffer	10182167	<b>Pass</b>

<b>Assay Name/Specification</b>	<b>Lot # 10184482</b>
<p><b>Blue-White Screening (Terminal Integrity)</b>            A sample of Litmus38i vector linearized with a 10-fold excess of EagI-HF<sup>™</sup>, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in &lt;1% white colonies.</p>	<b>Pass</b>
<p><b>Endonuclease Activity (Nicking)</b>            A 50 µl reaction in CutSmart<sup>™</sup> Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 20 Units of EagI-HF<sup>™</sup> incubated for 4 hours at 37°C results in &lt;20% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	<b>Pass</b>
<p><b>Exonuclease Activity (Radioactivity Release)</b>            A 50 µl reaction in CutSmart<sup>™</sup> Buffer containing 1 µg of a mixture of single and double-stranded [<sup>3</sup>H] E. coli DNA and a minimum of 100 units of EagI-HF<sup>™</sup> incubated for 4 hours at 37°C releases &lt;0.1% of the total radioactivity.</p>	<b>Pass</b>
<p><b>Ligation and Recutting (Terminal Integrity)</b>            After a 20-fold over-digestion of pXba DNA with EagI-HF<sup>™</sup>, &gt;95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,</p>	<b>Pass</b>

Assay Name/Specification	Lot # 10184482
<p>&gt;95% can be recut with Eagl-HF™.</p> <p><b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pXba DNA and a minimum of 100 Units of Eagl-HF™ incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p> <p><b>Protein Purity Assay (SDS-PAGE)</b> Eagl-HF™ is &gt;95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.</p>	<p style="text-align: center;"><b>Pass</b></p> <p style="text-align: center;"><b>Pass</b></p>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



YunJie Sun  
Production Scientist  
11 Apr 2023



Michael Tonello  
Packaging Quality Control Inspector  
13 Apr 2023