

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	DrallI-HF®
Catalog Number:	R3510S
Concentration:	20,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 μg of Lambda DNA in rCutSmart Buffer in 1 hour at 37°C in a total reaction volume of 50 μl.
Packaging Lot Number:	10240353
Expiration Date:	05/2026
Storage Temperature:	-20°C
Storage Conditions:	20 mM Tris-HCl, 300 mM NaCl, 1 mM Dithiothreitol, 0.1 mM EDTA, 200 μg/ml rAlbumin, 50% Glycerol (pH 7.4 @ 25°C)
Specification Version:	PS-R3510S/L/V v2.0

DrallI-HF® Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R3510SVIAL	DrallI-HF®	10240350	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10236230	Pass	
B6004SVIAL	rCutSmart™ Buffer	10237089	Pass	

Assay Name/Specification	Lot # 10240353
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in rCutSmart <sup>™</sup> Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 200 units of DraIII-HF® incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Functional Testing (15 minute Digest) A 50 μl reaction in rCutSmart™ Buffer containing 1 μg of Lambda DNA and 1 μl of DraIII-HF® incubated for 15 minutes at 37⁰C results in complete digestion as determined by agarose gel electrophoresis.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 10-fold over-digestion of Lambda DNA with DraIII-HF®, ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with DraIII-HF®.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of	Pass





240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10240353
20 units of DrallI-HF® incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	
<b>Protein Purity Assay (SDS-PAGE)</b> DraIII-HF® is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	Pass
<b>qPCR DNA Contamination (E. coli Genomic)</b> A minimum of 20 units of DraIII-HF® is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Ana Egana Production Scientist 17 May 2024

Michae

Michael Tonello Packaging Quality Control Inspector 17 May 2024

