

## New England Biolabs Certificate of Analysis

**Product Name:** BsiWI-HF<sup>®</sup>  
**Catalog Number:** R3553S  
**Concentration:** 20,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of PhiX174 DNA in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Packaging Lot Number:** 10076252  
**Expiration Date:** 11/2021  
**Storage Temperature:** -20°C  
**Storage Conditions:** 300 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 500 µg/ml BSA, (pH 7.4 @ 25°C)  
**Specification Version:** PS-R3553S/L v1.0

BsiWI-HF <sup>®</sup> Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R3553SVIAL	BsiWI-HF <sup>®</sup>	10060906	Pass
B7204SVIAL	CutSmart <sup>®</sup> Buffer	10074633	Pass
B7024SVIAL	Gel Loading Dye, Purple (6X)	10071081	Pass

Assay Name/Specification	Lot # 10076252
<b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in CutSmart <sup>®</sup> Buffer containing 1 µg of supercoiled pUC19 DNA and a minimum of 20 units of BsiWI-HF incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in CutSmart <sup>®</sup> Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 100 units of BsiWI-HF incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Functional Testing (15 minute Digest)</b> A 50 µl reaction in CutSmart <sup>®</sup> Buffer containing 1 µg of PhiX174 DNA and 1 µl of BsiWI-HF incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 20-fold over-digestion of PhiX174 DNA with BsiWI-HF, >95% of the DNA	Pass

Assay Name/Specification	Lot # 10076252
fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with BsiWI-HF.	
<b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in CutSmart® Buffer containing 1 µg of PhiX174 DNA and a minimum of 100 units of BsiWI-HF incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	<b>Pass</b>
<b>Protein Purity Assay (SDS-PAGE)</b> BsiWI-HF is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.



Penghua Zhang  
Production Scientist  
19 Jun 2020



Josh Hersey  
Packaging Quality Control Inspector  
19 Jun 2020