

New England Biolabs Product Specification

| | |
|-------------------------------|---|
| Product Name: | <i>NheI-HF</i> [®] |
| Catalog #: | R3131S/L |
| Concentration: | 20,000 units/ml |
| Unit Definition: | One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA (<i>HindIII</i> digest) in rCutSmart Buffer in 1 hour at 37°C in a total reaction volume of 50 µl. |
| Shelf Life: | 24 months |
| Storage Temp: | -20°C |
| Storage Conditions: | 250 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 0.15% Triton X-100, 200 µg/ml rAlbumin (pH 7.4 @ 25°C) |
| Specification Version: | PS-R3131S/L v2.0 |
| Effective Date: | 07 Nov 2023 |

Assay Name/Specification (minimum release criteria)

Blue-White Screening (Terminal Integrity) - A sample of LITMUS38i vector linearized with a 10-fold excess of *NheI-HF*[®], religated and transformed into an *E. coli* strain expressing the LacZ beta fragment gene results in <1% white colonies.

Endonuclease Activity (Nicking) - A 50 µl reaction in rCutSmart[™] Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 100 units of *NheI-HF*[®] incubated for 4 hours at 37°C results in <0.1% conversion to the nicked form as determined by agarose gel electrophoresis.

Exonuclease Activity (Radioactivity Release) - A 50 µl reaction in rCutSmart[™] Buffer containing 1 µg of a mixture of single and double-stranded [³H] *E. coli* DNA and a minimum of 200 units of *NheI-HF*[®] incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

Functional Testing (15 minute Digest) - A 50 µl reaction in rCutSmart[™] Buffer containing 1 µg of Lambda-HindIII DNA and 1 µl of *NheI-HF*[®] incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.

Ligation and Recutting (Terminal Integrity) - After a 100-fold over-digestion of Lambda HindIII DNA with *NheI-HF*[®], >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with *NheI-HF*[®].

Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in rCutSmart[™] Buffer containing 1 µg of Lambda-HindIII DNA and a minimum of 200 units of *NheI-HF*[®] incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Protein Purity Assay (SDS-PAGE) - *NheI-HF*[®] is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.



New England Biolabs Product Specification

| Assay Name/Specification (minimum release criteria) |
|---|
|---|

| |
|---|
| qPCR DNA Contamination (<i>E. coli</i> Genomic) - A minimum of 20 units of NheI-HF [®] is screened for the presence of <i>E. coli</i> genomic DNA using SYBR [®] Green qPCR with primers specific for the <i>E. coli</i> 16S rRNA locus. Results are quantified using a standard curve generated from purified <i>E. coli</i> genomic DNA. The measured level of <i>E. coli</i> genomic DNA contamination is ≤ 1 <i>E. coli</i> genome. |
|---|

One or more products referenced in this document may be covered by a 3rd-party trademark.
Please visit www.neb.com/trademarks for additional information.



Date 07 Nov 2023

Nancy Considine
Quality Approver

